

Parameterizing air-water gas exchange in the shallow, microtidal New River estuary

Bryce R. Van Dam ^{a,b,*}, James B. Edson ^c, Craig Tobias ^c

^a Present address: Department of Biological Sciences, Florida International University, Miami, FL 33199

^b Institute of Marine Sciences, University of North Carolina – Chapel Hill, NC 28557

^c Woods Hole Oceanographic Institution, Woods Hole, MA 02543

^d Department of Marine Sciences, University of Connecticut, Groton, CT 06340

Contents of this file

Figures S1 to S3




Figure S1. Footprint climatology for the New River estuary EC tower (using: Kljun et al., 2015), calculated from hourly QA/QC'd data. The black plus (+) depicts the location at which measurements were made. Footprint contour lines are shown in steps of 10 % from 10 to 90 %. The X- and Y-axis shows the distance in meters from the EC tower.




Figure S2. Box plots of average daily CO₂ flux (mmol m⁻² hr⁻¹) for the two representative days, 23 August 2017 (a) and 30 October 2017 (b), shown in figure 8. Differences were analyzed by one-way ANOVA, and a Tukey's least significant difference post-hoc procedure ($\alpha = 0.05$). Numbers indicate significant differences between group means. The lack of numbers in (a) indicates that the ANOVA revealed no significant effect of different k parameterizations on calculated CO₂ flux.




Figure S3. Relationship between convective velocity scale and k_{600} , showing linear fit and corresponding correlation coefficient (Rsq).